



Review

Discussion about Visual Dependence in Balance Control: European Society for Clinical Evaluation of Balance Disorders

Raphaël Maire, Arthur Mallinson, Hadrien Ceyte, Sebastien Caudron, Christian Van Nechel, Alexandre Bisdorff, Mans Magnusson, Hannes Petersen, Herman Kingma, Philippe Perrin

Clinic of Otolaryngology, Head and Neck Surgery, Neurotology Unit, Lausanne University Hospital, Lausanne, Switzerland (RM)

Neurotology Unit, Vancouver General Hospital, Vancouver, Canada (AM)

Division of Otolaryngology, University of British Columbia Faculty of Medicine, Vancouver, Canada (AM)

University of Lorraine, Development, Adaptation and Handicap, Faculty of Medicine and UFR STAPS, Villers-lès-Nancy, France (HC, SC, PP)

Unit of Neuro-Ophthalmology, Erasme University Hospital, Bruxelles, Belgium (CVN)

Department of Neurology, Emile Mayrisch Hospital, Esch-sur-Alzette, Luxembourg (AB)

Department of Otorhinolaryngology, Head and Neck Surgery, Clinical Sciences, University Hospital of Lund, Lund, Sweden (MM)

Department of Otorhinolaryngology, Landspítali University Hospital, Reykjavik, Iceland (HP)

Department of Otorhinolaryngology, Head and Neck Surgery, Clinical Sciences, Maastricht University Medical Center, Maastricht, The Netherlands (HK)

Faculty of Physics, Tomsk Research State University, Russia (HK)

Department of Pediatric Otolaryngology, University Hospital of Nancy, Vandoeuvre-lès-Nancy, France (PP)

Cite this article as: Maire R, Mallinson A, Ceyte H, Caudron S, Van Nechel C, Bisdorff A, et al. Discussion about Visual Dependence in Balance Control: European Society for Clinical Evaluation of Balance Disorders. J Int Adv Otol 2017; 13: 404-6.

The executive committee of the European Society for the clinical evaluation of balance disorders meets annually to address equilibrium problems that are not well understood. This is a review paper on discussions in the latest meeting we held. Materials and methods: Seeing patients with vestibular disorders who end up depending on visual information as part of their compensation process is a common clinical occurrence. However, this "visual dependence" can generate symptoms, which include nausea, sensations of imbalance, and anxiety. It is unclear how this develops, as symptoms can be widely variable from patient to patient. There are several triggering factors to this symptom set, and quantifying it in a given patient is extremely difficult. Results: The committee agreed that the presence of this symptom set can be suggestive of vestibular pathology, but the pathology does not have to be present. As a result, there is no correlation between symptom severity and test results. Conclusion: Visual dependence can often be present in a patient, although little, if any, measurable pathology is present. It is important to emphasize that although we cannot accurately measure this with either standardized testing or pertinent questionnaires, "hypersensitive" patients have a genuine disease and their symptoms are not of psychiatric origin.

KEYWORDS: Balance, visual dependence

INTRODUCTION

The ESCEBD has been meeting annually in Nancy, France, since 2005. It aims to discuss themes related to equilibrium that are not yet clearly defined or standardized ^[1, 2]. One of the latest discussions attempted to define and understand the concept of visual dependence and also attempted to develop parameters that could be used to diagnose this entity. The following is a brief report of our discussion.

BACKGROUND

Internal orientation in space and motion in humans are influenced by centrally integrated visual, vestibular, and proprioceptive inputs. On aging, the quality of all sensory inputs decreases. As a result, the elderly have a decreased ability to acquire and utilize each sensory input. Moreover, the senescence of somatosensory and vestibular information may make visual information more relevant (i.e., "preferred" input). Several studies have shown that aging is accompanied by an increased Romberg's quotient; this indicates that a greater reliance or weighting is being placed on visual information ^[3-5]. However, the onset of in-

Corresponding Address: Arthur Mallinson E-mail: art@mallinson.ca

Submitted: 27.07.2017 • Accepted: 08.08.2017

©Copyright 2017 by The European Academy of Otolology and Neurotology and The Politzer Society - Available online at www.advancedotology.org

creased visual dependence or reliance does not suggest pathology by itself. This increased visual dependence can put individuals at risk of developing many individual symptoms and limitations, particularly in the group of patients we regarded as having “vestibular hypersensitivity.” The presence of vestibular hypersensitivity is not necessarily a pathological situation, but it renders a patient more susceptible to becoming symptomatic from a small difference in visual and vestibular inputs.

Another confounding factor is that in the presence of vestibular pathology, increased reliance on visual information is part of the compensation process; this can generate symptoms ^[6].

DEFINITION OF VISUAL DEPENDENCE

The concept of “visual vertigo” was first described by Erasmus Darwin in 1797 ^[7]. Since the 1950s, the term “visual dependence” has been used to describe inappropriate reliance on visual cues in postural behavior in situations where it might be better to use somatosensory or vestibular inputs for balance. In the literature, various terminologies that include hypersensitivity (intolerance) to visual stimulations were proposed. “Phobic postural vertigo” is the term that was originally described by Brandt ^[8]. In 1995, Bronstein ^[9] suggested that it was the expression of enhanced visual dependence in patients with an undetected vestibular problem and coined the term “visual vertigo.” Other terms that are clinically used include “space motion discomfort” and “visual vestibular mismatch” ^[6]. The term “visually induced dizziness” has recently been introduced as an all-encompassing term ^[10], but mechanisms underlining the concept of visual dependence and how and why it develops are still poorly understood.

Visual dependence induces symptoms generated by static or moving visual cues in the environment (i.e., related to spatial orientation). These symptoms can be present alone or together with intolerance to passive/active body movements. Visual dependence can also result in signs, such as measurable imbalance, which are often described by patients as an “inherent sense of instability” ^[11]. Visual dependence is associated with other symptoms or part of syndromes, including vestibular diseases, anxiety, motion sickness, and migraine. It can also be an accompanying element to disorders such as agoraphobia, but ascribing these symptoms as having psychiatric origin should be avoided ^[12].

It is unclear how “visual dependence” develops, but we define it as “reduced ability to disregard visual cues in complex or conflicting visual environments (e.g., height vertigo, crowd, traffic, supermarkets, etc.)”. In brief, the reasons for the age-related development of visual dependence may correspond to the reduced ability to use visual inputs in a proper manner. Symptoms and signs can be provoked by an overabundance of visual cues in certain visual environments that was previously coped with.

FACTORS TRIGGERING VISUAL DEPENDENCE

Vestibular hypersensitivity by itself is an intolerance but instead is the inability to suppress stimuli that are environmentally inaccurate (e.g., moving train, 3D movies). The origin of visual dependence is unclear; symptoms can be generated by an assortment of pathological situations. The ESCEBD panel agrees that visual dependence is facilitated by the following:

1. vestibular end organ diseases, e.g., vestibular neuritis, that provoke an increase in visual contribution in the sensory integration process, often leading to difficulties in resolving various spatial daily situations where the whole body is implicated
2. vestibular migraine (which may be hypersensitivity) or conflicting visual input
3. psychiatric situations (i.e., environmental situations including panic and anxiety, which may be generated by stimuli such as watching the movement of water).
4. brain trauma (including blunt force trauma)

We feel that there are other non-pathological situations that can be implicated as triggering factors as visual dependence may also develop without any predisposing condition. In motion sickness (an inadequate response in certain individuals to environmental situations), vision is an important triggering factor and symptoms are facilitated in visually dependent subjects.

DISCUSSION

As outlined by several groups ^[6,9], visual dependence can suggest the presence of vestibular pathology. However, on aging, the level of visual dependence also increases ^[13]. It was initially suggested by Paige ^[14] that visual dependence resulted from a differing signal between two sensory inputs, visual senescence and adaptive plastic mechanisms, which normally maintain the vestibulo-ocular performance under conditions of head movement. If the senescence of these two systems did not occur in parallel, a resulting mismatch (and the possible development of visual dependence) in elderly patients might occur. Paige supported this by showing that the rate of caloric abnormalities was similar in elderly and younger patients with documented vestibular abnormalities. He suggested that the developed mismatch in both groups was similar (i.e., nonparallel senescence in elderly patients “mimicked” vestibular loss in younger patients). We agree that this decrease in the quality of sensory input is a factor that increases reliance on visual cues.

CLINICAL DIAGNOSIS AND ASSESSMENT OF VISUAL DEPENDENCE

As visual dependence creates symptoms and signs resulting from overweighting (or over-reliance) of visual inputs in the balance maintenance strategy, quantifying visual dependence should be possible using standardized assessments that employ visual stimuli such as optokinetic stimulation during standing eyes open, the rod-and-frame test, computerized dynamic posturography (e.g., conditions 3 and 6 of Equitest™ or similar testing on other equipment), or referenced visual stimuli in virtual reality conditions. However, currently, there is no correlation between symptoms and test results and only minimal correlation between the various testing methods reported in the literature. ESCEBD members involved in this discussion agree that no specific test for visual dependence exists to date and that this condition would be best evaluated using questionnaires.

There are three important points that we feel need to be emphasized:

1. Visual dependence frequently occurs in patients with a history of past vestibular disease but pathology does not have to be present as motion sickness and many other factors can generate these symptoms. A neurovestibular examination can display a past (inactive) peripheral vestibular deficit but is most often normal.

2. The symptoms of visual dependence can be generated by the actual compensation process (i.e., a conscious increase in reliance on visual stimuli can elicit symptoms).
3. The symptoms of visual dependence are often part of “chronic subjective dizziness” syndrome ^[15], which is a physical and not a psychiatric entity.

TREATMENT OF VISUAL DEPENDENCE

There is currently no standardized rehabilitation technique for visual dependence. Vestibular exercises are useful and help improve balance control. Optokinetic stimulation and virtual reality are tools that can often help, and it is thought this is accomplished by inducing symptoms and “habituating” a patient ^[16]. Both may be used as therapies, and both aim to lower visual sensitivity and restore central visuo-vestibular integration. Improvement in symptoms can be consolidated with a program of progressive confrontation to disabling situations. Learning from fighter pilots or rally car navigators might open new perspectives ^[17].

CONCLUSION

Visual dependence corresponds to hypersensitivity to moving or conflicting visual stimulations that can produce disabling vertigo/dizziness symptoms in triggering situations. Mechanisms underlining the symptomatology are unclear but likely imply the overweighting of vision in the equilibrium multisensory integration strategy. Symptoms frequently occur as part of syndromes including past vestibular disease and certain anxiety-related disorders, but the symptom set is not a psychiatric diagnosis or condition. In patients with visual dependence, an extensive history has to be obtained by a clinician familiar with the symptom set. These patients have a genuine disease, and their symptoms should not be regarded as having psychiatric origin. Standardized testing and pertinent questionnaires need to be developed to help us understand how to help such patients cope.

Peer-review: Externally peer-reviewed.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has received no financial support.

REFERENCES

1. Kingma H, Gauchard G, de Waele C, Van Nchel C, Bisdorff A, Yelnik A, et al. Stocktaking on the development of posturography for clinical use. *J Vest Res* 2011; 21: 117-25.
2. Maire R, Gauchard G, Deviterne D, Magnusson M, Kingma H, Perrin PP. European Society for Clinical Evaluation of Balance Disorders: discussion about dual-task conditions combining postural control with cognitive task. *Laryngoscope* 2010; 120: 2108-9. [\[CrossRef\]](#)
3. Lafont Ch, Dupui Ph, Costes-Salon MC. Equilibre et vieillissement. In: *L'année gérontologique*. Paris: Serdi, 1991: 151-91.
4. Pyykkö I, Aalto H, Hytonen M, et al. Effect of age on postural control. In: Amblard B, Berthoz A, Clarac F, eds. *Posture and gait: development, adaptation and modulation*. Amsterdam, New York, Oxford: Excerpta Medica 1988: 95-104.
5. Straube A, Botzel K, Hawken M, Paulus W, Brandt TH. Postural control in the elderly: differential effects of visual, vestibular and somatosensory input. In: Amblard B, Berthoz A, Clarac F, eds. *Posture and gait: development, adaptation and modulation*. Amsterdam, New York, Oxford: Excerpta Medica, 1988: 105-14.
6. Mallinson AI, Longridge NS. Visual vestibular mismatch in work-related vestibular injury. *Otol Neurotol* 2004; 26: 691-4.
7. Balaban CD, Jacob RG. Background and history of the interface between anxiety and vertigo. *J Anxiety Disord* 2001; 15: 27-51. [\[CrossRef\]](#)
8. Brandt TH. Phobic postural vertigo. *Neurology* 1996; 46: 1515-19. [\[CrossRef\]](#)
9. Bronstein AM. The visual vertigo syndrome. *Acta Otolaryngol Suppl* 1995; 520: 45-8. [\[CrossRef\]](#)
10. Bisdorff A, Von Brevern M, Lempert T, Newman-Toker DE. Classification of vestibular symptoms: Towards an international classification of vestibular disorders. *J Vestib Res* 2009; 1: 13.
11. Longridge NS, Mallinson AI. “Across the board” posturography abnormalities in vestibular injury. *Otol Neurotol* 2005; 26: 695-8. [\[CrossRef\]](#)
12. Furman JM, Jacob RG, Redfern MS. Clinical evidence that the vestibular system participates in autonomic control. *J Vest Res* 1998; 8: 27-34. [\[CrossRef\]](#)
13. Jamet M, Deviterne D, Gauchard GC, Vançon G, Perrin PP. Higher visual dependency increases balance control perturbation during cognitive task fulfilment in elderly people. *Neurosci Lett* 2004; 259: 61-4. [\[CrossRef\]](#)
14. Paige GD. Senescence of human visual-vestibular interactions. 1. Vestibulo-ocular reflex and adaptive plasticity with aging. *J Vest Res* 1992; 2: 133-51.
15. Staab JP. Chronic subjective dizziness. *Neuro-otology* 2012; 18: 1118-41. [\[CrossRef\]](#)
16. Pavlou M, Kanegaonkar RG, Swapp D, Bamiou DE, Slater M, Luxon LM. The effect of virtual reality on visual vertigo symptoms in patients with peripheral vestibular dysfunction: A pilot study. *J Vestib Res* 2012; 22: 273-81.
17. Perrin P, Lion A, Bosser G, Gauchard G, Meistelman C. Motion sickness in rally co-drivers, *Aviat Space and Environ Med* 2013; 84: 473-7. [\[CrossRef\]](#)